

## Math 1552: Integral Calculus

### Review Problems for Test 1, Sections 4.8, 5.1-5.6, 8.2-8.3

1. **Formula Recap:** complete each of the following formulas.

(a) The general Riemann Sum is found using the formula:

(b) Some helpful summation formulas are:

$$\sum_{i=1}^n c =$$

$$\sum_{i=1}^n i =$$

$$\sum_{i=1}^n i^2 =$$

(c) Properties of the definite integral:

$$\int_a^a f(x)dx =$$

$$\int_b^a f(x)dx =$$

$$\int_a^b cf(x)dx =$$

(d) State the Fundamental Theorem of Calculus:

(e) Using the FTC:

$$\frac{d}{dx} \left[ \int_{a(x)}^{b(x)} f(t)dt \right] =$$

(f) If  $F$  is an antiderivative of  $f$ , that means:

(g) If  $F$  is an antiderivative of  $f$ , then:

$$\int f(g(x))g'(x)dx =$$

$$\int_a^b f(g(x))g'(x)dx =$$

(h) To find the area between two curves, use the following steps:

(h) Evaluate an integral using *integration by parts* if:

To choose the value of  $u$ , use the rule: -----.

(i) To evaluate integrals with powers or products of trig functions, use the following trig identities to try to obtain a  $u$ -substitution:

2. Fill in the integration formulas below:

$$\int x^n dx, \quad (n \neq -1) =$$

$$\int \sin(ax) dx =$$

$$\int \cos(ax) dx =$$

$$\int \sec^2(ax) dx =$$

$$\int \sec(ax) \tan(ax) dx =$$

$$\int \csc(ax) \cot(ax) dx =$$

$$\int \csc^2(ax) dx =$$

$$\int \frac{1}{1 + (ax)^2} dx =$$

$$\int \frac{1}{\sqrt{1 - (ax)^2}} dx =$$

$$\int \frac{1}{x} dx =$$

$$\int e^{ax} dx =$$

$$\int b^{ax} dx =$$

$$\int \tan x dx =$$

$$\int \sec x dx =$$

$$\int \csc x dx =$$

$$\int \cot x dx =$$

Problems from Recitation Worksheets

1. True or False?

- (a) If  $F$  and  $G$  are both antiderivatives of  $f$ , then  $F = G$ .
- (b) The antiderivative of  $\sec^2(3x)$  is  $\frac{1}{3} \tan(3x)$ .
- (c) The indefinite integral of a function  $f$  is the collection of all antiderivatives of  $f$ .
- (d) We know how to find the antiderivative of  $\cos(x^2)$ , and it is  $\sin(x^2)$ .
- (e) To find the upper sum  $U_f$  of a function  $f$  on  $[a, b]$ , after partitioning the interval into  $n$  pieces, evaluate  $f$  at the right-hand endpoint of each subinterval.
- (f) When the interval  $[a, b]$  is partitioned into  $n$  pieces, there are exactly  $n$  endpoints.
- (g) A partition of the interval  $[a, b]$  does not need to be evenly spaced in order to calculate a Riemann Sum.
- (h) If  $f$  is positive and continuous on  $[a, b]$ , and  $A$  is the actual area bounded by  $f$ ,  $x = a$ ,  $x = b$ , and the  $x$ -axis, then  $L_f < A < U_f$ .
- (i) We always set  $x_i^*$  to be the right-hand endpoint of the  $i^{\text{th}}$  interval.
- (j)  $\sum_{i=1}^n i^2 = \left(\frac{n(n+1)}{2}\right)^2$ .
- (k) If  $f(x) \geq 0$  on  $[a, b]$ , then  $\int_a^b f(x) dx$  represents the total area bounded by  $f$ ,  $x = a$ ,  $x = b$ , and the  $x$ -axis.
- (l) If  $f$  is a continuous function, then the function  $F(x) = \int_a^x f(t)dt$  is an anti-derivative of  $f$ .
- (m) If  $F$  is an anti-derivative of  $f$ , then  $\int_a^b f(x)dx$  represents the slope of the secant line of  $F(x)$  on the interval  $[a, b]$ .
- (n)  $\frac{d}{dx} \left[ \int_a^b f(t)dt \right] = f(x)$ .
- (o) Given that  $f$  is continuous on  $[a, b]$  and  $F'(x) = f(x)$ , then  $F(b) - F(a)$  represents the net area bounded by the graph of  $y = f(x)$ , the lines  $x = a$ ,  $x = b$ , and the  $x$ -axis.
- (p)  $\int f(x)g(x) dx = \left(\int f(x) dx\right) \cdot \left(\int g(x) dx\right)$
- (q) To evaluate  $\int \sin^{-1}(x)dx$  by parts, choose  $u = \sin^{-1}(x)$  and  $dv = dx$ .
- (r) To evaluate  $\int x \ln(x) dx$  by parts, choose  $u = x$  and  $dv = \ln(x) dx$ .
- (s) To evaluate  $\int \cot(x) dx$ , integrate by substitution choosing  $u = \sin(x)$ .

2. Evaluate the following indefinite integrals.

(a)  $\int \left(\sqrt{x} - \frac{1}{x}\right)^2 dx$

(b)  $\int [4^{-2x} + e^{-5x}] dx$

(c)  $\int \left(\frac{e^{\sqrt{x}} + x^{\sqrt{x}}}{\sqrt{x}}\right) dx$

(d)  $\int \left(\frac{2}{3x} - \frac{1}{\sqrt{4-x^2}}\right) dx$

3. A particle travels with a velocity given by  $v(t) = -\frac{1}{3}t^2 + 4t + 2$ , where position is measured in meters and time in seconds.

(a) Find the acceleration of the particle when  $t = 1$  second.

(b) If the initial position is 4 m, find the position of the particle at  $t = 1$  second.

4. (*Applying the Riemann Sum*) You are driving when all of a sudden, you see traffic stopped in front of you. You slam the brakes to come to a stop. While your brakes are applied, the velocity of the car is measured, and you obtain the following measurements:

|                                  |    |    |    |    |    |   |
|----------------------------------|----|----|----|----|----|---|
| Time since applying breaks (sec) | 0  | 1  | 2  | 3  | 4  | 5 |
| Velocity of car (in ft/sec)      | 88 | 60 | 40 | 25 | 10 | 0 |

(a) Plot the points on a curve of velocity vs. time.

(b) Using the points given, determine upper and lower bounds for the total distance traveled before the car came to a stop.

5. Estimate the area under the graph of  $f(x) = 10 - x^2$  between the lines  $x = -3$  and  $x = 2$  using  $n = 5$  equally spaced subintervals, by finding:

(a) the upper sum,  $U_f$ .

(b) the lower sum,  $L_f$ .

6. (*Applying the Definite Integral*) A marketing company is trying a new campaign. The campaign lasts for three weeks, and during this time, the company finds that it gains customers as a function of time according to the formula:

$$C(t) = 5t - t^2,$$

where  $t$  is time in weeks and the number of customers is given in thousands.

Using the general form of the definite integral,

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \frac{b-a}{n} \sum_{i=1}^n f(x_i^*),$$

calculate the **average** number of customers gained during the three-week campaign.

7. Explain why the following property is true:

$$\left| \int_a^b f(x)dx \right| \leq \int_a^b |f(x)|dx.$$

Can you find an example where the inequality is strict?

8. Using the general form of the definite integral,  $\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*)\Delta x$ , evaluate:

$$\int_2^4 (x-1)^2 dx.$$

9. Evaluate  $\int_0^2 |x-1|dx$  using integral properties from class. (HINT: draw a picture, and use geometry!)

10. Suppose that  $f(x)$  is an even function such that  $\int_0^2 f(x)dx = 5$  and  $\int_0^3 f(x)dx = 8$ . Find the value of  $\int_{-2}^3 f(x)dx$ .

11. Evaluate the integrals:

(a)  $\int_1^2 \frac{3x-5}{x^3} dx$ .

(b)  $\int_2^5 (2-x)(x-5)dx$ .

(c)  $\int_{\pi}^{\frac{7\pi}{2}} \frac{1+\cos(2t)}{2} dt$ .

12. Find  $F'(2)$  for the function

$$F(x) = \int_{\frac{x}{2}}^{x^2} \left( \frac{t}{1-\sqrt{t}} \right) dt.$$

13. (a) Given the function  $f$  below, evaluate  $\int_1^9 f(x)dx$ .

$$f(x) = \begin{cases} x^2 + 4, & x < 4 \\ \sqrt{x} - x, & x \geq 4 \end{cases}$$

(b) Would you get the same answer to part (a) if you evaluated  $F(9) - F(1)$ ? What does this tell you about the FTC and continuity?

14. (a) Evaluate the expressions:

$$\int_0^1 x(1+x) dx, \quad \left( \int_0^1 x dx \cdot \int_0^1 (1+x) dx \right)$$

(b) Looking at your answer in part (a), what, if anything, can you say in general about  $\int (f(x) \cdot g(x)) dx$ ?

15. For each integral below, determine if we can evaluate the integral using the method of  $u$ -substitution. If the answer is "yes", evaluate the integral.

(a)  $\int \frac{1}{x^2} \sec\left(\frac{1}{x}\right) \tan\left(\frac{1}{x}\right) dx$

(b)  $\int x \csc^2(x) dx$

(c)  $\int \frac{\sin 3x - \cos 3x}{\sin 3x + \cos 3x} dx$

(d)  $\int \tan(x^2) dx$

16. Evaluate the following integrals using the method of substitution.

(a)  $\int \frac{1}{\ln(x^x)} dx$

(b)  $\int \frac{e^{2x}}{\sqrt{4-3e^{2x}}} dx$

(c)  $\int \frac{dx}{\sqrt{4-(x+3)^2}}$

17. Suppose that  $y = f(x)$  and  $y = g(x)$  are both continuous functions on the interval  $[a, b]$ . Determine if each statement below is always true or sometimes false.

(a) Suppose that  $f(c) > g(c)$  for some number  $c \in (a, b)$ . Then the area bounded by  $f$ ,  $g$ ,  $x = a$ , and  $x = b$  can be found by evaluating the integral  $\int_a^b (f(x) - g(x)) dx$ .

(b) If  $\int_a^b (f(x) - g(x)) dx$  evaluates to -5, then the area bounded by  $f$ ,  $g$ ,  $x = a$ , and  $x = b$  is 5.

(c) If  $f(x) > g(x)$  for every  $x \in [a, b]$ , then  $\int_a^b |f(x) - g(x)| dx = \int_a^b (f(x) - g(x)) dx$ .

18. Find the area bounded by the region between the curves  $f(x) = x^3 + 2x^2$  and  $g(x) = x^2 + 2x$ .

19. Find the area bounded by the region enclosed by the three curves  $y = x^3$ ,  $y = -x$ , and  $y = -1$ .

20. Find the area bounded by the curves  $y = \cos x$  and  $y = \sin(2x)$  on the interval  $[0, \frac{\pi}{2}]$ .

21. Find the area of the triangle with vertices at the points (0,1), (3,4), and (4,2). USE CALCULUS.

22. For each function below: (i) determine which method to use to evaluate the function (formula, u-substitution, or integration by parts, and (ii) evaluate the integral.

(a)  $\int_1^e \frac{\sqrt{\ln x}}{x} dx$

(b)  $\int (\ln x)^2 dx$

(c)  $\int x^2 e^{x^3} dx$

(d)  $\int x^3 e^{x^2} dx$

(e)  $\int 4^{-x} dx$

(f)  $\int x^2 \cdot 4^x dx$

23. Determine if each integral below can be evaluated using a method we have learned so far (formula, u-substitution, integration by parts, or trig identities). If so, evaluate the integral. If not, explain why it cannot be evaluated.

(a)  $\int x^5 \ln(x) dx$

(b)  $\int \sin^5(2x) \cos^3(2x) dx$

(c)  $\int \cos^2(3x) dx$

(d)  $\int \tan(x) \ln[\cos(x)] dx$

(e)  $\int \sin(x^2) dx$

(f)  $\int \tan^4(x) dx$

(g)  $\int e^{2x} \sin(3x) dx$



Additional Test Review Problems

24. True or false?

- (a) When evaluating a **definite** integral using  $u$ -substitution, different choices of  $u$  may lead to different final answers.
- (b) Integration by Parts is a Product Rule in integral form.
- (c) The goal of integration by parts is to go from an integral  $\int f'(x)g'(x)dx$  that we can't evaluate to an integral  $\int f(x)g(x)dx$  that we can evaluate.
- (d) Definite integrals can not be evaluated by Integration by Parts.
- (e) If  $f$  is a continuous, increasing function, then the right-hand Riemann sum method always overestimates the definite integral.
- (f) Let  $f$  be a continuous function and  $av(f)$  be the average of  $f$ . Then  $av(f) \cdot (b - a) = \int_a^b f(x)dx$ .
- (g) When finding the area between the curves  $y = x^3 - x$  and  $y = x^2 + x$  it suffices to find the value of the definite integral  $\int_{-1}^2 [(x^3 - x) - (x^2 + x)] dx$ , and then take the absolute value of this value to get the right answer.
- (h) To find the area between the curves  $y = x^3 - x$  and  $y = x^2 + x$ , first set the equations equal and solve to find the intersection points  $x = -1$  and  $x = 2$ , plug in a test-point into the equations or graph the curves to determine **top** and **bot**, and then evaluate  $\int_{-1}^2 (\mathbf{top}) - (\mathbf{bot}) dx$ .
- (i) If  $\int_0^1 f(x) dx = 9$  and  $f(x) \geq 0$ , then  $\int_0^1 \sqrt{f(x)} dx = 3$ .

25. Evaluate the following integrals.

- (a)  $\int_0^{\frac{\pi}{4}} \sec^2(t)e^{1+\tan(t)} dt$
- (b)  $\int \sin^3(x) \cos^3(x) dx$
- (c)  $\int \frac{1}{\sqrt{4-9w^2}} dw$
- (d)  $\int x \sin(x) \cos(x) dx$
- (e)  $\int \sec^4(x) dx$
- (f)  $\int \ln(x + 1) dx$

26. Suppose:  $f(1) = 2, f(4) = 7, f'(1) = 5, f'(4) = 3$  and  $f''(x)$  is continuous. Find the

value of:

$$\int_1^4 x f''(x) dx.$$

27. Consider the following limit

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \cos\left(\pi \cdot \frac{i}{n}\right) \cdot \frac{\pi}{2n}.$$

- (a) Express the limit as a definite integral.  
(b) Compute the definite integral from part (a).

28. Let  $f(x) = 3x + 4$ .

- (a) Estimate the area of the region between the graph of  $f$ , the lines  $x = -1$  and  $x = 2$ , and the  $x$ -axis using an upper sum with three rectangles of equal width.  
(b) Find the actual area in part (a) by taking the limit of a general Riemann Sum using  $n$  equally spaced subintervals, and taking  $x_i^*$  as the right-hand endpoint of each interval.

29. Find the area bounded by the curves  $y = \cos^2(x)$  and  $y = -\sin^2(x)$ , and the lines  $x = 0$  and  $x = \pi$ . (Hint: draw a picture in GeoGebra - an online graphing tool.)

30. Find the area bounded by the curves  $y = -x^2 + 6x$  and  $y = x^2 - 2x - 24$ . (Hint: sketch the curves or make a sign chart.)

31. Find  $F'(4)$  if

$$F(x) = \int_{\frac{x^2}{4}}^{x^2} \ln(\sqrt{t}) dt.$$

32. What value of  $b > -1$  maximizes the integral:

$$\int_{-1}^b x^2(7-x)dx?$$

33. Find a number  $c$  so that  $f(c)$  is equal to the average value of the function  $f(x) = 1 + x$  on the interval  $[-1, 3]$ . Graphically, what does that mean?

## Answers

- (b), (c), (g), (k), (l), (o), (q), (s) are true
- (a)  $\frac{1}{2}x^2 - 4\sqrt{x} - \frac{1}{x} + C$   
(b)  $-\frac{1}{2\ln 4}4^{-2x} - \frac{1}{5}e^{-5x} + C$   
(c)  $2e^{\sqrt{2}}\sqrt{x} + \frac{1}{\sqrt{2}+1/2}x^{\sqrt{2}+1/2} + C$   
(d)  $\frac{2}{3}\ln|x| - \sin^{-1}\left(\frac{x}{2}\right) + C$
- (a)  $\frac{10}{3} m/s^2$ , (b)  $7\frac{8}{9} m$
- (b) Upper: 223 ft, Lower: 135 ft
- (a) 44 (b) 31
- 4,500 customers
- Consider the difference between NET and TOTAL area.
- $\frac{26}{3}$
- 1
- 13
- (a)  $-\frac{3}{8}$ ; (b)  $\frac{9}{2}$ ; (c)  $\frac{5\pi}{4}$
- 24
- (a)  $\frac{79}{6}$ ; (b) you cannot use the FTC as stated when  $f$  is discontinuous somewhere on the interval  $[a, b]$
- (a)  $\frac{5}{6}$  and  $\frac{3}{4}$ ; no general rule
- (a)  $-\sec\left(\frac{1}{x}\right) + C$ , (c)  $-\frac{1}{3}\ln|\sin 3x + \cos 3x| + C$
- (a)  $\ln|\ln x| + C$ , (b)  $-\frac{1}{3}\sqrt{4 - 3e^{2x}} + C$ , (c)  $\sin^{-1}\left(\frac{x+3}{2}\right) + C$
- (c) is true
- $\frac{37}{12}$
- $\frac{5}{4}$
- $\frac{1}{2}$
5. 4.5
- (a)  $\frac{2}{3}$   
(b)  $x(\ln x)^2 - 2x \ln x + 2x + C$   
(c)  $\frac{1}{3}e^{x^3} + C$

- (d)  $\frac{x^2 e^{x^2}}{2} - \frac{e^{x^2}}{2} + C$
- (e)  $-\frac{1}{\ln 4} 4^{-x} + C$
- (f)  $\frac{1}{\ln 4} x^2 \cdot 4^x - \frac{2}{(\ln 4)^2} x \cdot 4^x + \frac{2}{(\ln 4)^3} 4^x + C$
23. (a)  $\frac{x^6 \ln x}{6} - \frac{x^6}{36} + C$
- (b)  $\frac{1}{12} \sin^6(2x) - \frac{1}{16} \sin^8(2x) + C$
- (c)  $\frac{1}{2}x + \frac{1}{12} \sin(6x) + C$
- (d)  $-\frac{1}{2}(\ln[\cos(x)])^2 + C$
- (e) Cannot be evaluated
- (f)  $\frac{1}{3} \tan^3(x) - \tan(x) + x + C$
- (g)  $\frac{2}{13} e^{2x} \sin(3x) - \frac{3}{13} e^{2x} \cos(3x) + C$
24. (e), (f) are true
25. (a)  $e^2 - e$
- (b)  $\frac{1}{6} \cos^6(x) - \frac{1}{4} \cos^4(x) + C$
- (c)  $\frac{1}{3} \arcsin\left(\frac{3w}{2}\right) + C$
- (d)  $\frac{x}{2} \sin^2 x - \frac{1}{4}x + \frac{1}{8} \sin 2x + C$
- (e)  $\tan(x) + \frac{\tan^3(x)}{3} + C$
- (f)  $(x+1) \ln(x+1) - (x+1) + C$
26. 2
27. (a)  $\int_0^{\frac{\pi}{2}} \cos(2x) dx$ , (b) 0
28. (a) 21, (b) 16.5
29.  $\pi$
30.  $\frac{512}{3}$  or approximately 170.67
31.  $14 \ln 2$
32.  $b = 7$
33.  $c = 1$