

**MATH 2550/2551 READING DAY STUDY SESSION WORKSHEET**

Problems marked with \*\* are *only* relevant for MATH 2551.

PROBLEMS

1. Find  $T$ ,  $N$ , and curvature for  $r(t) = (3 \sin t)\vec{i} + (3 \cos t)\vec{j} + 4t\vec{k}$ .
2. \*\*Write acceleration in terms of tangential and normal components for  $r(t) = (t + 1)\vec{i} + 2t\vec{j} + t^2\vec{k}$ ,  $t = 1$ .
3. Find the limit:  $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2 - y^2 + 5}{x^2 + y^2 + 2}$ .
4. Find all second-order partial derivatives for  $w = x \sin(x^2y)$ .
5. Evaluate  $\frac{dw}{dt}$  for  $w = 2ye^x - \ln z$ ,  $x = \ln(t^2 + 1)$ ,  $y = \arctan t$ ,  $z = e^t$ ,  $t = 1$ .
6. Find  $\frac{\partial z}{\partial x}$ ,  $\frac{\partial z}{\partial y}$  for  $z^3 - xy + yz + y^3 - 2 = 0$ ,  $(1, 1, 1)$ .
7. Find the derivative of the function at the point in the direction of  $\vec{v}$  for  $f(x, y) = 2xy - 3y^2$ ,  $P = (5, 5)$ ,  $\vec{v} = 4\vec{i} + 3\vec{j}$ .
8. Find the tangent plane and normal line for  $2z - x^2 = 0$ ,  $P_0(2, 0, 2)$ .
9. Find all local maxima, minima, and saddle points for  $f(x, y) = x^2 + xy + y^2 + 3x - 3y + 4$ .
10. Find absolute maxima and minima for  $f(x, y) = 2x^2 - 4x + y^2 - 4y + 1$  on the closed triangular plate bounded by the lines  $x = 0$ ,  $y = 2$ ,  $y = 2x$  in the first quadrant.
11. Find the maximum value of  $f(x, y) = 49 - x^2 - y^2$  on the line  $x + 3y = 10$ .
12. Write an iterated integral using (a) vertical cross-sections and (b) horizontal cross-sections for  $y = \sqrt{x}$ ,  $y = 0$ ,  $x = 9$ .
13. Sketch the region and evaluate  $\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy$ .
14. Change to polar and evaluate  $\int_{-1}^1 \int_0^{\sqrt{1-x^2}} dy dx$ .
15. For the region below  $z = 4 - xy$  and in  $0 \leq x \leq 2$ ,  $0 \leq y \leq 1$ , integrate the function  $3 - 4x$ .
16. Use  $x = \frac{u}{v}$ ,  $y = uv$  in  $\mathbb{R}$  (first quadrant) bounded by  $xy = 1$ ,  $xy = 9$ ,  $y = x$ ,  $y = 4x$  for  $\iint \left( \sqrt{\frac{y}{x}} + \sqrt{xy} \right) dx dy$ .
17. \*\*Evaluate the line integral  $\int (xy + y + z) ds$  along the curve  $r(t) = 2t\vec{i} + t\vec{j} + (2 - 2t)\vec{k}$ ,  $0 \leq t \leq 1$ .
18. \*\*Find the flow of the field  $F = -4xy\vec{i} + y8\vec{j} + 2\vec{k}$  along  $r(t) = t\vec{i} + t^2\vec{j} + \vec{k}$ ,  $0 \leq t \leq 2$ .
19. \*\*Find the circulation and flux of the field  $F = x\vec{i} + y\vec{j}$  around  $r(t) = \cos t\vec{i} + \sin t\vec{j}$ ,  $0 \leq t \leq 2\pi$ .
20. \*\*Find the potential function for  $F = e^{y+2z}(\vec{i} + x\vec{j} + 2x\vec{k})$ .

21. \*\*Use Green's Theorem to find counterclockwise circulation and outward flux for field  $F = (y^2 - x^2)\vec{i} + (x^2 + y^2)\vec{j}$  and curve  $C: y = 0, x = 3, y = x$ .
22. \*\*Use a parametrization to express the area of the surface as a double integral: the portion of the cone  $z = 2\sqrt{x^2 + y^2}$  between the planes  $z = 2$  and  $z = 6$ .
23. Evaluate  $\iint_S 2y \, dV$  over  $S: y^2 + z^2 = 4$  between  $x = 0, x = 3 - z$ .
24. At time  $t = 0$ , a particle is located at the point  $(0, 1, 2)$ . At this time it is traveling towards the point  $(3, 2, 3)$ , has speed 3 at  $(0, 1, 2)$ , and has constant acceleration  $3\vec{i} - \vec{j} + \vec{k}$ . Find an equation for the position vector  $\vec{r}(t)$  at time  $t$ .
25. Let  $P(A_0, B_0, C_0), Q(A_1, B_1, C_1)$  be two distinct points in 3-dimensional space.
  - (a) Write down a vector parameterization for the line  $PQ$ .
  - (b) Show that the curvature  $\kappa$  of  $PQ$  is 0.
  - (c) \*\*Show that the torsion  $\tau$  of  $PQ$  is 0.
26. (a) At what points  $(x, y)$  in the plane is the function  $f(x, y) = \frac{y}{1 + \cos x}$  continuous?  
 (b) Find the following limit by first rewriting the fraction  $\lim_{(x,y) \rightarrow (1,-1)} \frac{x^3 + y^3}{x + y}$ .
27. Let  $C$  be the smooth curve given by the intersection of the two surfaces  $xyz = 1$  and  $x^2 + 2y^2 + 3z^2 = 6$ .
  - (a) Write down an equation that describes  $C$  implicitly.
  - (b) Write down a parametric equation for the line tangent to  $C$  at the point  $(1, 1, 1)$ .
28. Let  $D$  be the cylinder bounded below by  $z = -1$ , bounded on the sides by  $x^2 + y^2 = 1$ , and bounded above by  $z = 1$ . (**Do not evaluate the integrals.**)
  - (a) Express the volume of  $D$  as an iterated triple integral in cylindrical coordinates.
  - (b) Express the volume of  $D$  as an iterated triple integral in rectangular coordinates.
  - (c) Express the volume of  $D$  as an iterated triple integral in spherical coordinates.
29. Consider the function  $f(x, y) = \frac{\ln(x^2 + y^2)}{\sqrt{x^2 + y^2}}$ .
  - (a) Describe the region  $R$  given by  $1 \leq x^2 + y^2 \leq e$  in polar coordinates.
  - (b) Write down the integral of  $f(x, y)$  over  $R$  in polar coordinates.
  - (c) Evaluate the integral you found in part (b).
30. Let  $D$  be the region in  $xyz$ -space defined by the inequalities  $1 \leq x \leq 2, 0 \leq xy \leq 2, 0 \leq z \leq 1$ . Consider the coordinate transformation of  $D$  to the  $uvw$ -plane given by  $u = x, v = xy, w = 3z$ .
  - (a) Sketch the preimage  $G$  of  $D$  under the coordinate transformation in the  $uvw$ -plane and label the bounding curves.
  - (b) Write down the Jacobian associated to this coordinate transformation.
  - (c) Evaluate the integral  $\iiint_D (x^2y + 3xyz) \, dx \, dy \, dz$ .
31. A flat circular plate has the shape of the region  $x^2 + y^2 \leq 1$ . Points on the plate have temperature  $T(x, y) = x^2 + 2y^2 - x$ . Find the temperatures of the hottest and coldest points of the plate.
32. Find the points on the surface  $xyz = 1$  closest to the origin.
33. Find the volume of the wedge cut from the cylinder  $x^2 + y^2 \leq 1$  by the planes  $z = -y$  and  $z = 0$ .

## ANSWERS

1.  $T = \frac{3\cos t}{5}\vec{i} - \frac{3\sin t}{5}\vec{j} + \frac{4}{5}\vec{k}$ ,  $N = (-\sin t)\vec{i} - (\cos t)\vec{j}$ ,  $\kappa = \frac{3}{25}$
2.  $a(1) = \frac{4}{3}T + \frac{2\sqrt{5}}{3}N$
3.  $\frac{5}{2}$
4.  $w_{xx} = 6xy \cos(x^2y) - 4x^3y^2 \sin(x^2y)$ ,  $w_{yy} = -x^5 \sin(x^2y)$ ,  $w_{xy} = 3x^2 \cos(x^2y) - 2x^4y \sin(x^2y)$
5.  $\frac{dw}{dt} = 4t \arctan t + 1 = \pi + 1$
6.  $\frac{1}{4}, -\frac{3}{4}$
7.  $-4$
8.  $2x - z - 2 = 0$ ,  $r(t) = (2, 0, 2) + t(-4, 0, 2)$
9.  $f(-3, 3) = -5$  minimum
10.  $(0, 0) = 1$ , abs max,  $(1, 2) = -5$ , abs min
11. 39
12.  $0 \leq x \leq 9$ ,  $0 \leq y\sqrt{x}$ ,  $0 \leq y \leq 3$ ,  $y^2 \leq x \leq 9$
13.  $e - 2$
14.  $\frac{\pi}{2}$
15.  $\int_0^2 \int_0^1 \int_0^{4-xy} (3 - 4x) dz dy dx = -\frac{17}{3}$
16.  $\int_1^2 \int_1^3 \frac{(u+v)2u}{v} du dv = 8 + \frac{52}{3} \ln 2$
17.  $\frac{13}{2}$
18. 48
19. Circulation = 0; Flux =  $2\pi$
20.  $f(x, y, z) = xe^{y+2z} + C$
21. Flux =  $-9$ , Circulation = 9
22.  $\int_0^{2\pi} \int_1^3 r\sqrt{5} dr d\theta$
23.  $r(u, v) = r(x, \theta) = \langle x, 2 \sin \theta, 2 \cos \theta \rangle$   
 $\iint f(r(u, v)) |r_u \times r_v| dA = \int_0^{2\pi} \int_0^{3-2\cos \theta} 2(2 \sin \theta)(2) dx d\theta = 0$
24.  $\vec{r}(t) = \left(\frac{3}{2}t^2 + \frac{9}{\sqrt{11}}t\right)\vec{i} + \left(-\frac{1}{2}t^2 + \frac{3}{\sqrt{11}}t + 1\right)\vec{j} + \left(\frac{1}{2}t^2 + \frac{3}{\sqrt{11}}t + 2\right)\vec{k}$
25. (a)  $L(t) = \langle x(t) = A_0 + (A_1 - A_0)t, y(t) = B_0 + (B_1 - B_0)t, z(t) = C_0 + (C_1 - C_0)t \rangle$

MATH 2550/2551 - Spring 2018

26. (a)  $f$  is continuous on  $\{(x, y) | x \neq (2k + 1)\pi; k = \dots, -1, 0, 1, \dots\}$ . (b) 1
27. (a)  $x^2 + 2y^2 + 3z^2 = 6xyz$ . (b)  $L(t) = \langle 1 + 2t, 1 - 4t, 1 + 2t \rangle$
28. (a)  $\int_{-1}^1 \int_0^{2\pi} \int_0^1 r \, dr \, d\theta \, dz$ . (b)  $\int_{-1}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \int_{-1}^1 dz \, dx \, dy$   
 (c)  $\int_0^{2\pi} \int_{\pi/4}^{3\pi/4} \int_0^{1/\sin \varphi} \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta + 2 \int_0^{2\pi} \int_0^{\pi/4} \int_0^{1/\cos \varphi} \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$
29. (a)  $1 \leq r^2 \leq e$ . (b)  $\int_0^{2\pi} \int_1^{\sqrt{e}} 2 \ln(r) \, dr \, d\theta$ . (c)  $-2\pi\sqrt{e} + 4\pi$
30. (b)  $\frac{1}{3u}$ . (c)  $2 + 3 \ln(2)$
31. Hottest:  $\frac{9}{4}$  at points  $(-\frac{1}{2}, \frac{\sqrt{3}}{2})$ ,  $(-\frac{1}{2}, -\frac{\sqrt{3}}{2})$ . Coldest:  $-\frac{1}{4}$  at point  $(\frac{1}{2}, 0)$ .
32. Points  $(1, 1, 1)$ ,  $(-1, -1, -1)$ ,  $(1, -1, -1)$ , and  $(-1, 1, -1)$  are all at distance 3 from the origin.
33.  $\frac{2}{3}$